Product Data Sheet



Product Name 3-D Life Scrambled RGD Peptide

Catalog Number P11-3

Description

RGD sequences are motifs on extracellular matrix proteins promoting adhesion to cellular integrin receptors. The *3-D Life* Scrambled RGD Peptide contains a scrambled RGD motif and a thiol group on a N-terminus. The thiol group is used to immobilize the peptide on maleimide polymers in *3-D Life* Hydrogels. The immobilized peptide is used in control experiments to determine the specificity of cell adhesion caused by a genuine RGD peptide (e.g. *3-D Life* RGD Peptide).

For instructions, please consult General Protocols GP-1 "Preparation of 3-D Life Fast Gelling Hydrogels" or GP-2 "Preparation of 3-D Life Slow Gelling Hydrogels" and the 3-D Life Hydrogels User Guide on www.cellendes.com.

Quantity

3x 1 µmol

Components

	Material	Quantity	Concentration of reactive groups	Storage
	Scrambled RGD peptide, lyophilized	3x 1 µmol	20 mmol/L*	Lyophilisate and after reconstitution: -20°C or lower
0	Water	600 µl	n/a	Room temperature or lower

All materials are filter-sterilized.

Reconstitution

Scrambled RGD Peptide:

- Briefly centrifuge vial containing the Scrambled RGD Peptide lyophilisate to make sure that entire product is at the bottom of the reaction tube.
- 2. Add 48 µl *3-D Life* Water per tube of Scrambled RGD Pepetide to dissolve lyophilisate at a concentration of 20 mmol/L thiol groups.
- 3. Close tube and briefly vortex.
- 4. Incubate for 5 min.
- 5. Briefly vortex and centrifuge again.
- 6. The Scrambled RGD Peptide is now ready for use.

Pept. Sequence Acetyl-Cys-Doa*-Doa-Gly-Arg-Asp-Gly-Ser-Pro-NH2

*: Doa:8-amino-3,6-dioxaoctanoic acid

Molar Mass 1022 g/mol

Purity >90% (HPLC)

Intended for research use only. Not for use in human therapeutic or diagnostic applications.

Cellendes GmbH Gerhard-Kindler-Str. 8, 72770 Reutlingen, Germany Tel.: +49 7121 15940 0

www.cellendes.com

^{*}Volume/concentration after reconstitution of lyophilisate.